




PCT/EP200 4 / 0 1 3 2 5 2



INVESTOR IN PEOPLE

The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ




I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed


Dated 2 November 2004

BEST AVAILABLE COPY

**PRIORITY  
DOCUMENT**

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

Patents Form 1/77

Patents Act 1977  
(Rule 16)The  
Patent  
Office

1/77

## Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office  
Cardiff Road  
Newport  
Gwent NP9 1RH

6 OCT 2004

1. Your reference	GB Case BT/3-22351/P2			0422070.3	
2. Patent application (The Patent Office will...)					
3. Full name, address and postcode of the or of each applicant (underline all surnames)	Ciba Specialty Chemicals Water Treatments Limited Cleckheaton Road Low Moor Bradford West Yorkshire BD12 0JZ 7885391005 England				
Patent ADP number (if you know it)					
If the applicant is a corporate body, give the country/state of its incorporation	England				
4. Title of invention	MICROORGANISM				
5. Name of your agent (if you have one)	Ciba Specialty Chemicals Water Treatments Limited				
"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)	Patents Department PO Box 38 Cleckheaton Road Low Moor Bradford West Yorkshire BD12 0JZ 07685391002				
Patents ADP number (if you know it)					
6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number	Country	Priority application number (if you know it)	Date of filing (day/month/year)		
7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application	Number of earlier application		Date of filing (day/month/year)		
8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if: a) any applicant named in part 3 is not an inventor, or b) there is an inventor who is not named as an applicant, or c) any named applicant is a corporate body. (see note (d))	YES				

Patents Form 1/77

## Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form -

Description **24** /

Claim(s) **4** /

Abstract **1** /

Drawing(s) -

10. If you are also filing any of the following, state how many against each item.

Priority documents -

Translations of priority documents -

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*) -

Request for preliminary examination and search (*Patents Form 9/77*) **1** /

Request for substantive examination (*Patents Form 10/77*) -

Any other documents -  
(please specify)

11. I/We request the grant of a patent on the basis of this application

Signature

Date

**06 October 2004**

*C. Choppen*

12. Name and daytime telephone number of person to contact in the United Kingdom

**Catherine Choppen**

**01274 417446**

## Warning

*After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.*

## Notes

- a) *If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.*
- b) *Write your answers in capital letters using black ink or you may type them.*
- c) *If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.*
- d) *Once you have filled in the form you must remember to sign and date it.*
- e) *For details of the fee and ways to pay please contact the Patent Office.*

Patents Form 1/77

BT/3-22351/P2

1

Microorganism

The present invention relates to a microorganism and to methods of storing the microorganism. The invention also relates to a novel nitrile hydratase enzyme  
5 and also to a method of converting a nitrile to an amide employing the nitrile hydratase enzyme.

It is well known to employ biocatalysts, such as microorganisms that contain enzymes, for conducting chemical reactions. Nitrile hydratase enzymes are  
10 known to catalyse the hydration of nitriles directly to the corresponding amides. Typically nitrile hydratase enzymes can be produced by a variety of microorganisms, for instance microorganisms of the genus Bacillus, Bacteridium, Micrococcus, Brevibacterium, Corynebacterium, Pseudomonas, Acinetobacter, Xanthobacter, Streptomyces, Rhizobium, Klebsiella,  
15 Enterobacter, Erwinia, Aeromonas, Citrobacter, Achromobacter, Agrobacterium, Pseudonocardia and Rhodococcus.

Many references have described the synthesis of nitrile hydratase within microorganisms. Arnaud et al., Agric. Biol. Chem. 41: (11) 2183-2191 (1977)  
20 describes the characteristics of an enzyme they refer to as 'acetonitrilase' in Brevibacterium sp R312 which degrades acetonitrile to acetate via the amide intermediate. Asano et al., Agric. Biol. Chem. 46: (5) 1183-1189 (1982) isolated Pseudomonas chlororaphis B23 which produced nitrile hydratase to catalyse the conversion of acrylonitrile to acrylamide, generating 400 g/L acrylamide. The  
25 article by Yamada et al., Agric. Biol. Chem. 50: (11) 2859-2865 (1986) entitled, "Optimum culture conditions for production by Pseudomonas chlororaphis B23 of nitrile hydratase", considered the optimisation of the medium components of the growth medium, including the inducer added for nitrile hydratase synthesis. Methacrylamide was found to be the best inducer for this organism.  
30 Methacrylamide was included in the culture at the start of growth.

Various strains of the *Rhodococcus rhodochrous* species have been found to very effectively produce nitrile hydratase enzyme.

EP-0 307 926 describes the culturing of *Rhodococcus rhodochrous*, specifically strain J1 in a culture medium that contains cobalt ions. A process is described for biologically producing an amide in which a nitrile is hydrated by the action of a nitrile hydratase produced by *Rhodococcus rhodochrous* J1, which has been cultured in the presence of cobalt ion. The use of various inducers (including crotonamide) is described for the synthesis of nitrile hydratase. In one embodiment an amide is produced in a culture medium of the microorganism in which a substrate nitrile is present. In another embodiment a substrate nitrile is added to the culture medium in which a nitrile hydratase has been accumulated to conduct the hydration reaction. There is also a description of isolating the microbial cells and supporting them in a suitable carrier, for instance by immobilisation, and then contacting them with a substrate. The nitrile hydratase can be used to hydrate nitriles into amides, and in particular the conversion of 3-cyanopyridine to nicotinamide.

EP-0 362 829 describes a method for cultivating bacteria of the species *Rhodococcus rhodochrous* comprising at least one of urea and cobalt ion for preparing the cells of *Rhodococcus rhodochrous* having nitrile hydratase activity. Specifically described is the induction of nitrile hydratase in *Rhodococcus rhodochrous* J1 using urea or urea derivatives which markedly increases the nitrile hydratase activity. Urea or its derivatives are added to the culture medium in one batch at a time or sequentially and cultivation occurs over 30 hours or longer, for instance up to 120 hours.

An article by Nagasawa et al., Appl. Microbiol. Biotechnol. 34: 783-788 (1991), entitled "Optimum culture conditions for the production of cobalt-containing nitrile hydratase by *Rhodococcus rhodochrous* J1", describes isolating J1 as an acetonitrile utilising strain which synthesises two different nitrile hydratases and

a nitrilase depending upon the culture conditions used. One nitrile hydratase is induced optimally by urea and urea analogues. Urea is added at the start of the culturing process and seems to become efficient as an inducer only when the basal medium is nutrient rich. Induction of the enzyme started gradually and  
5 increased in growth until it reached a maximum after 5 days of cultivation. The activity was found to decrease on prolonged cultivation.

Rhodococcus rhodochrous J1, is also used commercially to manufacture acrylamide monomer from acrylonitrile and this process has been described by  
10 Nagasawa and Yamada Pure Appl. Chem. 67: 1241-1256 (1995).

Leonova et al., Appl. Biochem. Biotechnol. 88: 231-241 (2000) entitled, "Nitrile Hydratase of Rhodococcus", describes the growth and synthesis of nitrile hydratase in Rhodococcus rhodochrous M8. The nitrile hydratase synthesis of  
15 this strain is induced by urea in the medium, the urea also acting as a nitrogen source for growth by this organism. Cobalt is also required for high nitrile hydratase activity. This literature paper looks at induction and metabolic effects in the main.

20 Leonova et al., Appl. Biochem. Biotechnol. 88: 231-241 (2000) also states that acrylamide is produced commercially in Russia using Rhodococcus rhodochrous M8. Russian patent 1731814 describes Rhodococcus rhodochrous strain M8.

25 Rhodococcus rhodochrous strain M33 that produces nitrile hydratase without the need of an inducer such as urea is described in US-A-5827699. This strain of microorganism is a derivative of Rhodococcus rhodochrous M8.

The production of acrylamide monomer in particular is desirable via the  
30 biocatalytic route. In the review publication by Yamada and Kobayashi, Biosci. Biotech. Biochem. 60: (9) 1391-1400 (1996) titled "Nitrile Hydratase and its

Application to Industrial Production of Acrylamide" a detailed account of the development of a biocatalytic route to acrylamide is described. Three successively better catalysts and their characteristics for acrylamide production and in particular the third generation catalyst *Rhodococcus rhodochrous* J1 are described in some detail.

A major disadvantage with the use of biocatalysts is the general lack of stability observed with wet microbial material during storage, transportation and use. Even with relatively stable enzymes and bacteria such as nitrile hydratases in *Rhodococcal* cells, the potential for spoilage before use has led to acceptance within the industry for the need to process the biocatalyst cell suspension in some way e.g. by freezing or freeze-drying of the aqueous mixture or alternatively immobilisation of the cells in some polymer matrix. In order to achieve maximum productivity from the biocatalyst it is important that the maximum biocatalytic activity is retained during its preparation and storage prior to use. In Chaplin and Bucke (1990) In: Enzyme Technology, published by Cambridge University Press, p 47 (Enzyme preparation and use) it was recognised that enzyme inactivation can be caused by heat, proteolysis, sub optimal pH, oxidation denaturants and irreversible inhibitors. A number of substances may cause a reduction in the rate of an enzymes ability to catalyse a reaction. This includes substances that are non-specific protein denaturants, such as urea.

In the presentation, Protein Stability, by Willem JH van Berkel, Wageningen University, factors that may cause deactivation or unfolding were considered and these included proteases, oxidation due to the presence of oxygen or oxygen radicals and denaturing agents causing reversible unfolding, such as urea.

Chaplin and Bucke (1990) In Enzyme Technology, published by Cambridge University Press, p 73 (Enzyme preparation and use) revealed that the key

factor regarding the preservation of enzyme activity involves maintaining the conformation of the enzyme structure. Therefore it was considering important to prevent unfolding, aggregation and changes in the covalent structure. Three approaches for achieving this were considered: (1) use of additives; (2) the  
5 controlled use of covalent modification; and (3) enzyme immobilisation.

EP-B-0 243 967 describes the preservation of nitrile hydration activity of nitrilase by the addition of stabilizing compounds selected from nitriles, amides and organic acids and their salts, to a solution or suspension of the enzyme or the  
10 immobilized form of the enzyme. It clearly states in the description that while a solution or suspension of a microorganism capable of producing nitrilase that hydrates nitriles such as acrylonitrile, to produce the corresponding amides such as acrylamide may be stored at room temperature as long as the storage period is short, storage at a low temperature, especially at a temperature in the  
15 vicinity of 0°C is preferred. It was described in EP-A-0 707 061 that addition of inorganic salts at a concentration of between 100 mM to the saturation concentration of the inorganic salts to an aqueous medium containing either a suspension of microbial cells or immobilized microbial cells, preserved the cells and enzyme activity for a prolonged period of time. This technique is described  
20 for the preservation of microbial cells that have nitrile hydratase or nitrilase activity. The addition of bicarbonate or carbonate salts to an aqueous solution of immobilized or unimmobilised microbial cells having nitrilase activity is described in US-B-6,368,804. Immobilisation has frequently involved removal of the enzyme from the whole cell, before immobilising the enzyme in a matrix.  
25 However, although such immobilisation provides very good protection for the enzyme, extraction of the enzyme from the whole cell is an intricate step, which can be time-consuming, expensive and can lead to loss of enzyme. Additionally whole microbial cells can be immobilized. US-A-5,567,608 provides a process of immobilising whole cell biocatalyst in a cationic copolymer which has good  
30 storage stability and prevents putrefaction.

Rhodococcus rhodochrous J1, which is used commercially to manufacture acrylamide monomer, is immobilised to (a) allow transportation and (b) to increase the longevity of the biocatalyst in use. In US-A-5,567,608 the inventors state that biocatalysts are normally immobilized for use on an industrial scale, to facilitate ease of separation of the biocatalyst from the reaction product, preventing impurities from the biocatalyst eluting into the product and to assist in continuous processes and recycling of the biocatalyst. However, immobilisation is an extra processing step that requires an additional plant and the use of potentially a number of other raw materials such as alginate, carrageenan, acrylamide and other acrylate monomers, and vinyl alcohol. Thus, this is an expensive processing step.

Various other ways have been proposed for minimising the deleterious effects of enzyme inactivation in an attempt reduce the negative impact on a chemical reaction process.

It is also known to freeze dry biocatalysts in order to preserve the activity of an enzyme in storage over a prolonged period of time. Again this is a potentially expensive processing step that is normally carried out with biocatalysts prepared on a small scale. Cryopreservation in liquid nitrogen or in the vapour phase of liquid nitrogen also affords long-term storage of microbial cells but requires a constant supply of liquid nitrogen. Freezing of recovered biomass or semi-pure or pure enzymes at temperatures of  $<-18^{\circ}\text{C}$  is also known to preserve biocatalytic activity for prolonged periods of time.

Furthermore, once the cell mass is introduced to the reactor and the reaction is taking place minimisation of the loss of efficacy is critical to the operational efficiency and the process economics. Once again, immobilisation of the microbial cells into some polymer matrix is standard procedure to optimise these process parameters.

It would therefore be desirable to provide a process and a biocatalyst where these disadvantages can be overcome.

According to the present invention we provide a microorganism that is

5 Rhodococcus rhodochrous strain NCIMB 41164 or a mutant thereof.

This new microorganism has been found to readily produce nitrile hydratase.

We have found that this new microorganism (and the nitrile hydratase produced therefrom) can be used in a process of converting nitriles, to the amide.

10 Rhodococcus rhodochrous NCIMB 41164 is particularly of use for the conversion of (meth)acrylonitrile to (meth)acrylamide. The microorganism and enzyme have been found to remain active, and in some cases even increase in activity, over long periods of time and furthermore can be recovered from the reaction mixture with undiminished activity after preparation of acrylamide at  
15 >50% w/w. Thus it can be, if required, reused either directly or after a further period of storage.

The details of the new strain Rhodococcus rhodochrous NCIMB 41164 are given below:

20

#### 1. Origin and Deposition

The Rhodococcus rhodochrous strain was isolated by us from soil in Bradford, England and deposited on 5<sup>th</sup> March 2003 at the National Collection of Industrial and Marine Bacteria (NCIMB), where it was assigned the accession number

25 NCIMB 41164 under the Budapest Treaty.

#### 2. Taxonomic identification of the microorganism

Identification of the soil isolate was carried out using the technique of 16S rDNA analysis. The sequence of the 16S rDNA gene obtained from the soil isolate  
30 was compared with nucleic acid sequence databases. The sequence obtained was compared to those found in a proprietary database (Microseq™) and the

top 20 hits were determined. Comparison of the sequence with this database identified the best match as *Rhodococcus rhodochrous* with a 97.48% similarity. This is a genus level match, but was most likely to be a strain of *Rhodococcus rhodochrous*. A further search against the public EMBL database  
5 identified the best match for this database to *Rhodococcus rhodochrous* with 99.698% similarity.

### 3. Morphological and cultural characteristics

- (1) Polymorphic growth
- 10 (2) Motility: immotile
- (3) Non-spore former
- (4) Gram positive
- (5) Aerobic
- (6) Growth on nutrient agar gives salmon pink round colonies within 48  
15 hours at 30°C

### 4. Cultivation and Nitrile Hydratase Synthesis

The *Rhodococcus rhodochrous* NCIMB 41164 of the present invention can be cultured under any conditions suitable for the purpose in accordance with any of  
20 the known methods, for instance as described in the aforementioned prior art. Preferably the microorganism is cultured in a culture medium that comprises urea or a derivative of urea. We have found that this microorganism can be grown in a medium containing acetonitrile or acrylonitrile as an inducer of the nitrile hydratase. In the presence of urea or urea derivative as an inducer and  
25 cobalt chloride as a source of cobalt ions, very high nitrile hydratase activity is achieved. For example urea and cobalt are added to the medium described in the experimental examples.

Desirably the *Rhodococcus rhodochrous* NCIMB 41164 can be cultured to give  
30 high enzyme activity, for instance about 250-300,000  $\mu\text{mol min}^{-1}/\text{g}$  dry biomass at 15°C. High nitrile hydratase activity can be achieved if urea or a urea

derivative is present in the culture medium. It may be present at the start of the culture or it may be added at some point during growth, but generally should be added before the onset of the stationary phase of growth. High nitrile hydratase activity can preferably be achieved if urea or the urea derivative is not present in  
5 any substantial amount in the culture medium at the start of the microorganism growth but is introduced later. By this we mean that urea or the urea derivative is not present or is present in an amount of less than 0.2 g/l, preferably less than 0.1 g/l. More preferably the culture medium is substantially free (i.e less than 0.2 g/l) of urea or the urea derivative for at least the first six hours of  
10 microorganism growth. It is especially preferred if the growth medium of the microorganism is substantially free of urea or the urea derivative for at least 12 hours and in some cases at least 24 hours before the introduction of the urea or the urea derivative as the growth rate of the microorganism is higher in the absence of urea or the urea derivative, but that it is added before 48 hours  
15 culturing of the microorganism. We have found that this enables higher nitrile hydratase activity to occur in a shorter period of time than if the urea or the urea derivative had been added at the start of culturing.

The invention also relates to a nitrile hydratase obtainable from a  
20 microorganism which is *Rhodococcus rhodochrous* NCIMB 41164 or a mutant thereof.

A further aspect of the invention concerns a process of preparing an amide from the corresponding nitrile wherein the nitrile is subjected to a hydration reaction  
25 in an aqueous medium in the presence of a biocatalyst selected from the group consisting of a microorganism which is *Rhodococcus rhodochrous* NCIMB 41164, a mutant thereof and a nitrile hydratase obtainable from *Rhodococcus rhodochrous* NCIMB 41164 or a mutant thereof. Hereafter the term 'biocatalyst' refers to the nitrile hydratase that is synthesised within the *Rhodococcus*  
30 *rhodochrous* NCIMB 41164 cell and may include the *Rhodococcus rhodochrous* NCIMB 41164 cell itself. Thus, the biocatalyst could be used as a whole cell

10

preparation in a fermentation medium, as an aqueous suspension, as a recovered cell paste as an immobilized cell preparation or as any other form of the nitrile hydratase suitable for the conversion of nitrile to amide that satisfies the requirements of this invention.

5

This process is particularly suitable for readily preparing an amide from the corresponding nitrile. In particular aqueous solutions of amide can be prepared in high concentration. The process is especially suitable for preparing acrylamide or methacrylamide.

10

The biocatalyst may be used as a whole cell catalyst for the generation of amide from nitrile. It may be immobilised for instance entrapped in a gel or it may be used preferably as a free cell suspension. Alternatively the nitrile hydratase enzyme may be extracted and for instance used directly in the process of preparing the amide.

15

In one preferred way of carrying out the process the biocatalyst is introduced into an aqueous medium suitable for carrying out the culturing of the microorganism. Typically a suspension of the biocatalyst, for instance whole cells of the microorganism, may be formed. A nitrile, for instance acrylonitrile or methacrylonitrile is fed into the aqueous medium comprising the biocatalyst in such a way that the concentration of (meth) acrylonitrile in the aqueous medium is maintained at up to 6% by weight. Nitrile such as acrylonitrile or methacrylonitrile is more preferably fed into the reaction medium and the reaction allowed to continue until the concentration of amide, for instance acrylamide or methacrylamide reaches the desired level, in particular between 30 and 55% by weight. Most preferably the concentration is around 50% by weight.

20

25

30

This new strain of *Rhodococcus rhodochrous* (NCIMB 41164) is capable of producing aqueous acrylamide solutions in high concentration (for instance 50%

acrylamide). Desirably the reaction may be carried out as a free cell process using a fed-batch type reactor to which the biocatalyst (*Rhodococcus rhodochrous* NCIMB 41164) is added in the form of fermentation broth or as harvested biomass.

5

The activity of the biocatalyst (*Rhodococcus rhodochrous* NCIMB 41164) and the nitrile hydratase produced therefrom is such that it can be recycled and reused for further hydration of nitrile to the corresponding amide.

- 10 Recycling of the biocatalyst is particularly suitable for any case of converting (meth) acrylonitrile to (meth) acrylamide. Thus in the manufacture of acrylamide when the reaction process is complete and acrylamide has been produced at the appropriate concentration, the catalyst can be removed and re-used to produce another batch of acrylamide without loss in nitrile hydratase activity.
- 15 This can even be achieved after the biocatalyst has been stored in water for several days (for instance three days) prior to reuse. It is even possible to prepare a third batch of acrylamide, even after further storage.

- According to one aspect of the invention we provide an aqueous composition
- 20 comprising a biocatalyst that is or is obtainable from the microorganism *Rhodococcus rhodochrous* strain NCIMB 41164 or a mutant thereof and wherein the biocatalyst is in the form of a non-actively growing free cell microorganism. We also provide a method of storing the biocatalyst, that is in the form of a non-actively growing free cell microorganism.

25

- The microbial cells of the biocatalyst used to carry out the conversion of nitrile to amide, may be regarded as a non-actively growing culture. By this we mean that the medium and the storage conditions in which the microorganism is held would not be expected to promote growth. The storage medium can for
- 30 instance be the *Rhodococcus rhodochrous* NCIMB 41164 cells that maybe recovered from the fermentation medium. Or the cells maybe used directly in

the fermentation medium, or they may be present as an aqueous suspension in a suitable suspending medium for instance; water; physiological saline solution; a suitable buffer solution such as phosphate buffer or any other similar buffer or a growth medium where metabolism in the microorganism cells is substantially zero as determined by measuring the growth rate, or the biomass concentration or oxygen consumption or nutrient consumption, or other form of measurement generally used to monitor microbial growth and metabolism.

The composition of the storage medium may comprise any residual fermentation broth components. The fermentation broth may include any of the typical ingredients used for culturing the microorganism and also may include products and by-products produced by the microorganism. Typical components of the fermentation broth include sugars, polysaccharides, proteins, peptides, amino acids, nitrogen sources, inorganic salts, vitamins, growth regulators and enzyme inducers. Specifically this could include monosaccharides or disaccharides as sugars; ammonium salts or other nitrogen sources; inorganic salts such as phosphates, sulphates, magnesium, calcium, sodium and potassium salts; metal compounds; vitamins; and complex fermentation medium components, for example corn steep liquor, peptone; yeast extract; organic or inorganic compounds that may be used for specific microbial growth requirements; specific enzyme inducers (such as urea that is used to induce the nitrile hydratase of *Rhodococcus rhodochrous* NCIMB 41164); and organic acids such as citrate or pyruvate; and any other organic or inorganic compounds that may be required to ensure successful growth of the *Rhodococcus rhodochrous* NCIMB 41164.

Usually when a biocatalyst, such as one that produces nitrile hydratase, is stored without continued growth for a period of time, even for a few days, it is normal to remove the microbial cells from the fermentation broth, whether it is the cells that are required as the catalyst, or whether the enzyme is recovered from the the cells or fermentation medium. This is to prevent microbial growth

in the fermentation broth causing putrefaction of the broth and to reduce protease activity that can cause the breakdown of the enzyme that is required. It is normal therefore to preserve the fermentation broth *per se* or to remove the cells to prevent the degradation of the biocatalyst through extraneous biological activity such as microbial contamination. The biocatalytic activity could normally be expected to reduce in a very short period of time such as within a day and certainly in less than two days if this were not carried out.

Methods of preserving the activity during the storage of biocatalysts, even for periods of time up to one-week, have normally involved removal of the biocatalyst from the fermentation broth and/or immobilisation of the biocatalyst in a suitable matrix and/or stabilisation using stabilising substances which then either become contaminants in the reaction mixture and this may be a problem further downstream or an additional processing step is required to remove the stabilizing compound or additive from the microbial cell suspension before it is used as a biocatalyst.

In the absence of such preservation treatments and normally biocatalysts that are kept at ambient temperatures tend to lose activity to the extent that they are no longer as effective or even suitable for catalysing reactions.

Growth of a microorganism for use as a biocatalyst may take place over a period of several days. During this time the microorganism is actively growing, that is to say balanced growth where the biomass is increasing together with an increase in and maintenance of the overall chemical composition of the cell.

Normally the growth of microorganisms is limited either by the exhaustion of nutrient or the accumulation of toxic products of metabolism and the growth rate reduces. Growth is maintained by feeding appropriate nutrients and maintaining a correct temperature and pH for growth and where required supplying oxygen.

The storage method described here promotes effective stability such that the biocatalyst can be readily used without any significant loss in activity. Storage stability is achieved without the necessity of resorting to for instance immobilisation, addition of stabilizing compounds or freeze drying. Storage  
5 stability may be achieved without resorting to removal of any of the fermentation broth components such as urea or urea derivatives, even though urea is a known protein deactivator.

The composition or the environment used in the method of storage may contain  
10 oxygen or can be a substantially oxygen free environment. By oxygen free we mean that the concentration of oxygen should be less than 1% dissolved oxygen concentration. Removal of oxygen from the fermentation broth can be achieved by any of the conventional methods for removing oxygen. These include purging for a period of time with an inert gas, removal of any head-  
15 space in the storage container, storing under diminished pressure or the addition of known oxygen scavengers such as ascorbic acid or hydrazine and hydrazide.

It would have been expected that after 2 days and especially after several days  
20 storage there would be some loss in nitrile hydratase activity. This would have been expected even in the absence of oxygen. It would have been expected especially in the presence of residual fermentation broth components, such as urea, and also at temperatures of above 0°C. This is because protease enzymes in the biocatalyst might be expected to break down other proteins in  
25 the cell, including the nitrile hydratase. Furthermore, the presence of urea or urea derivative could be expected to be detrimental, since urea is known to be a protein deactivator. However, the biocatalyst suffers none of the expected disadvantages and thus suffers no significant loss in nitrile hydratase activity.

30 On the contrary we find that during the storage period the activity of the biocatalyst comprising nitrile hydratase can in some cases actually increase.

Thus in another aspect of the invention we provide a method of increasing the nitrile hydratase activity of a biocatalyst capable of forming nitrile hydratase by storing the biocatalyst in a storage medium in accordance with the storage method of the present invention. Therefore, the method can result in a new  
5 biocatalyst composition by virtue of its increased activity. Therefore, nitrile hydratase of the biocatalyst composition, and in particular formed during storage of the biocatalyst, is new. Also, the biocatalyst does not produce the mal odours associated with putrefaction during the storage period.

10 Preferably the storage method allows the biocatalyst to be stored for at least two days and more preferably one or more weeks. In particular the biocatalyst may be stored from three to twenty eight days, for example 3 to 14 days.

The presence of fermentation broth components such as urea are not essential  
15 to the composition or the storage method of this aspect of invention. Where fermentation broth components are present, this may be urea or a urea derivative. The urea derivative can be for example an alkyl derivative of urea.

Urea or the urea derivative could be present in the biocatalyst composition  
20 through its inclusion in the fermentation mixture. In one form of the invention the composition or storage medium containing the biocatalyst may be deoxygenated and contain fermentation broth components such as urea.

A particularly advantageous feature of this aspect of the invention is that it is no  
25 longer necessary to separate the biocatalyst from the fermentation mixture in which it was cultured. This is of significant value since it avoids the requirement for an additional processing step. Therefore the composition may also comprise a fermentation mixture, which is then stored. In the method of storing the biocatalyst, we find that this may also be achieved in the presence of a  
30 fermentation mixture without any detrimental effects on the activity of the enzyme. This then allows the fermentation broth to be used immediately to

catalyse the reaction, or to allow it to be stored for several days or even weeks without detriment whilst the bioconversion step is being carried out also over a period of several days, thus ensuring a constant supply of readily available biocatalyst without need for additional processing steps thus simplifying and  
5 reducing the cost of the bioconversion step.

The biocatalyst may conveniently be stored at temperatures above its freezing point. Typically the biocatalyst may be stored at ambient temperatures, for instance up to 30 or 40°C. However, the advantage of the present method is  
10 that the biocatalyst may be stored at ambient temperatures without any special precautions for monitoring and controlling the temperature. Preferably the biocatalyst is stored at a temperature between 4 and 30 or 40°C, more preferably between 5 and 25°C, such as between 10 and 25°C and in particular 15 to 25°C.

15

According to a further aspect of the present invention we provide a method of producing an amide by contacting the corresponding nitrile by a nitrile hydratase,

in which the biocatalyst is part of a composition or stored in the form of a non-  
20 actively growing free cell microorganism in a storage medium in which the composition or storage medium comprises fermentation broth, and the biocatalyst is (or is obtainable from) the microorganism *Rhodococcus rhodochrous* strain NCIMB 41164 or a mutant thereof.

25 Thus in accordance with this aspect of the invention the biocatalyst may have been held in an environment containing oxygen or held in an oxygen-free environment. It may or may not contain residual fermentation broth components such as urea prior to commencing the conversion of the nitrile. This may be resulting from storing the biocatalyst in accordance with the storage aspect of  
30 the present invention or alternatively provided as a composition in accordance with the present invention.

As given previously it is not necessary to remove the biocatalyst from the fermentation mixture in which the biocatalyst has been prepared. Thus in a preferred form the environment in which the biocatalyst is held also contains components of a fermentation broth. Therefore a biocatalyst composition  
5 containing components of a fermentation broth can be combined with a nitrile which is then hydrated to the corresponding amide. We have found surprisingly that in contrast to previous knowledge, for instance in US-A-5,567,608 states that immobilisation of the biocatalyst is preferable to prevent elution of impurities from the biocatalyst into the reaction product, that the inclusion of fermentation  
10 broth in the reaction mixture does not affect the quality of the final product and this aspect is described in our co-filed UK application 0327901.5, identified by case number BT/3-22349/P1.

The fermentation mixture will comprise essential components for allowing  
15 microorganisms to be grown and sustained. In general the mixture will at least contain a carbon source, nitrogen source and various nutrients. This may include a saccharide for instance a monosaccharide such as glucose or other sugar or a disaccharide or polysaccharide, ammonium salts, complex medium components such as yeast extract and peptone, amino acids, vitamins,  
20 phosphate salts, potassium, sodium, magnesium and calcium salts, trace elements such as iron, cobalt, manganese, copper, zinc and the like. These and other ingredients can be included in the fermentation mixture at concentrations suitable for the particular microorganism. It is known that fermentations can be subject to changes in the productivity of the biocatalyst and the fermentation  
25 broth may be used at different stages of growth and so it is important to be able to store the biocatalyst after production in such a way.

We find that the activity of the biocatalyst does not diminish significantly during the reaction for a prolonged period. Consequently the biocatalyst may be  
30 replaced less frequently. Preferably the biocatalyst is used for a period of at least 2 days and loses substantially no activity over that period.

Generally the catalysis of the reaction using nitrile hydratase enables the nitrile to be converted into the corresponding amide in a single step. This process is of particular value when the nitrile is acrylonitrile and the amide is acrylamide. It is desirable to carry out this conversion step several times using a single batch of biocatalyst from which portions are removed over a period of several days to carry out several reactions where nitrile is converted to amide. Thus, it is important to be able to store the biocatalyst as inexpensively as possible without detriment to the catalyst whilst the bioconversion step is carried out simultaneously. So in effect one batch of biocatalyst can be stored ready for use to make several batches of for instance acrylamide. Several batches could be from 5 to 10 or more batches, even 15 to 20 batches.

In a further aspect of the invention we have found a way of improving the biocatalytic activity of a microorganism. The microorganism would be cultured in a culture medium that comprises urea or a derivative of urea. However, urea or the derivative of urea is introduced into the culture medium at least six hours after the start of growth of the microorganism. Normally the culture medium is substantially free of urea or the urea derivative for at least the first six hours of culturing the microorganism and thereafter urea or a urea derivative is added to the culture medium. As indicated previously by substantially free we mean that the culture medium contain less than 0.2 g/l, usually less than 0.1 g/l and may contain no urea or the urea derivative. Preferably the culture medium is substantially free of urea or the urea derivative for at least 12 hours and sometimes at least 24 hours. However, in order to maximise the biocatalytic activity it is preferred to introduce the urea or the urea derivative within 48 hours of culturing.

The biocatalytic activity can be established in terms of enzyme activity as described herein.

Preferably the microorganism is capable of producing a nitrile hydratase. Suitably a biocatalyst comprising such a microorganism can be used to prepare amides from the corresponding nitrile by a hydration process in which nitrile hydratase catalyses the reaction. The culturing of the microorganism by  
5 delayed introduction of urea or urea derivative provides increased nitrile hydratase activity particularly suitable for this reaction. The process is particularly suitable for the preparation of (meth) acrylamide from (meth) acrylonitrile. Such a process may be carried out as described herein. In addition the biocatalyst may be recycled and reused.

10

It is particularly desirable that the microorganism is of the *Rhodococcus* genus, preferably a *Rhodococcus rhodochrous* species, especially *Rhodococcus rhodochrous* NCIMB 41164.

15 The following examples provide an illustration of how to carry out the invention.

Example 1

Rhodococcus rhodochrous NCIMB 41164 was isolated from soil using an enrichment culture technique and it was grown on a medium containing the following constituents (g/l):  $\text{KH}_2\text{PO}_4$ , 7.0;  $\text{KH}_2\text{PO}_4$ , 3.0; peptone, 5.0; yeast  
5 extract, 3.0; glucose, 5.0;  $\text{MgSO}_4$ , 0.5; trace metals solution, 5 ml; acetonitrile, 20 ml. The pH was adjusted to 7.2. The nitrile hydratase activity was 4,000  $\mu\text{mol min}^{-1}/\text{g}$  dry cells at 15°C after 3 days growth at 28°C.

Example 2

10 (1) Rhodococcus rhodochrous NCIMB 41164 was grown in a 2L baffled Erlenmeyer flask containing 400 mL culture medium containing the following constituents (g/L): diPotassium hydrogen phosphate 0.7; Potassium hydrogen phosphate 0.3; glucose 10.0; peptone, 1.0; yeast extract 3.0; magnesium sulphate heptahydrate 0.5; Urea 5.0; cobalt chloride hexahydrate 0.01; tap  
15 water to 1L. The pH of the medium was adjusted to pH 7.2. The culture was grown at 28°C for 5 days after which the nitrile hydratase activity was 47,900  $\mu\text{mol min}^{-1}/\text{g}$  at 15°C.

20 (2) (a) Rhodococcus rhodochrous NCIMB 41164 was grown in the medium described in (1) except that peptone was omitted.

(b) Rhodococcus rhodochrous NCIMB 41164 was grown in the medium described in (2a) except that peptone was omitted as was urea. The organism was cultured for 24 hours and then 5 g/L urea was added to the culture which was grown for a further 5 days.

25 (C) Rhodococcus rhodochrous NCIMB 41164 was grown in the medium described in (2a) except that urea was not included in the medium. The organism was cultured for 48 hours and then 5 g/L urea was added to the culture which was grown for a further 4 days.

30 (d) Rhodococcus rhodochrous NCIMB 41164 was grown in the medium described in (2a) except that urea was not included in the medium. The organism was cultured for 6 days.

Samples were taken from the four cultures described above at time = 1, 2, 3 and 6 days after growth commenced. The nitrile hydratase activities were measured at 15°C, see table 1.

5 Table 1

Urea addition time (days)	Nitrile Hydratase Activity $\mu\text{mol min}^{-1}/\text{mg dry cells}$			
	T = 1 day	T = 2 days	T = 3 days	T = 6 days
0	9.1	24.2	24.8	37.6
1	1.0	21.6	49.3	41.3
2	ND	ND	15.1	15.3
None added	0.94	ND	0.46	0.98

ND not determined

### Example 3

- (1) *Rhodococcus rhodochrous* NCIMB 41164 was grown in a 280L fermenter containing 180 L culture medium containing the following constituents (g/L): diPotassium hydrogen phosphate 0.7; Potassium hydrogen phosphate 0.3; glucose 2.0; yeast extract 3.0; magnesium sulphate heptahydrate 0.5; cobalt chloride hexahydrate 0.01;. The pH of the medium was adjusted to pH 7.2. The culture was grown at 30°C for 3 days. Urea was added to the culture after 17 h. The nitrile hydratase activity was measured (at 30°C) periodically. 22 h after the urea was added the activity was approximately 176,000  $\mu\text{mol min}^{-1}/\text{g}$  at 30°C and after a further 9 h the activity had increased to 323, 000  $\mu\text{mol min}^{-1}/\text{g}$ .
- (2) 625 g of water was charged to the reactor to which *Rhodococcus rhodochrous* NCIMB 41164 was added. The mixture was heated to 25°C. Acrylonitrile 375 g was fed to the reactor at a rate to maintain the concentration

at 2% (w/w). After 175 minutes all of the acrylonitrile had been converted to acrylamide to a final concentration of approximately 50% (w/w).

(3) The cells from 2 were recovered by centrifugation and they were suspended in 625 g water. This suspension was stored at 4°C for 3 days prior to re-charging to the reactor. The procedure described in 5 was followed and again after 175 minutes all of the acrylonitrile was converted to acrylamide.

(4) The cells from 3 were treated as described in 3 above except they were stored for 2 days prior to re-use. Again 50% acrylamide was synthesised. The acrylic acid concentrations measured for the batches of acrylamide generated in example 3 2-4) (5-7 are shown in Table 2.

Table 2: Acrylic acid concentrations measured in each of the acrylamide batches

Example number	Acrylic Acid Concentration (ppm)
3-2	5650
3-3	102
3-4	None detected (<10 ppm)

#### 15 Example 4

(1) *Rhodococcus rhodochrous* NCIMB 41164 was grown in a 280L fermenter containing 180 L culture medium containing the following constituents (g/L): diPotassium hydrogen phosphate 0.7; Potassium hydrogen phosphate 0.3; glucose 1.0; yeast extract 3.0; magnesium sulphate heptahydrate 0.5; cobalt chloride hexahydrate 0.01; urea, 5.0. The pH of the medium was adjusted to pH 7.2. The culture was grown at 30°C for 3 days.

25L of the fermentation broth was degassed with nitrogen for 20 minutes prior to storage at ambient temperature, which was approx. 5°C for 3½ days. The nitrile hydratase activity was measured 15 h after harvesting and it was found to be 242,000 U/g at 25°C. When the NH activity was re-measured 3 days later it was found to be 293,000 U/g.

**Example 5**

Rhodococcus rhodochrous NCIMB 41164 was grown in a 2 L Erlenmeyer flask for 5 days at 28°C with shaking at 180 rpm in a culture medium containing the following constituents in g/L: diPotassium hydrogen phosphate 0.7; Potassium hydrogen phosphate 0.3; glucose 10.0; yeast extract 3.0; urea 5.0; magnesium sulphate heptahydrate 0.5; cobalt chloride hexahydrate 0.01;. The pH of the medium was adjusted to pH 7.2. The culture broth was divided into two portions, one half of which was deoxygenated using nitrogen. Portions of both the deoxygenated and the oxygenated culture broth were incubated at 4, 15 and 25°C for 1 week. The nitrile hydratase activity of the portions was measured periodically.

The results of the nitrile hydratase assays are shown in Table 3. The results are given in U/mg dry cells

**Table 3**

Incubation temp.	Time (days)					
	0	1	2	3	5	7
4°C (O <sub>2</sub> )	140	286		232	267	257
4°C (degassed)		274			214	293
15°C (O <sub>2</sub> )						
15°C (degassed)	140	218				
25°C (O <sub>2</sub> )	140	143				
25°C (degassed)		154	230			

It can be seen from the results in Example 5 that the biocatalyst can be stored effectively at ambient temperatures. Furthermore it can be seen that the nitrile

hydratase activity increased on this occasion on storage in comparison to day zero.

Example 6

- 5 Defrosted cells of *Rhodococcus rhodochrous* NCIMB 41164 were resuspended in water. The nitrile hydratase activity was measured over a period of 1 week. The relative nitrile hydratase activities measured are shown in Table 4.

Table 4

Time (days)	Relative nitrile hydratase activity (%)		
	4°C	15°C	25°C
0	100	100	100
1	66	64	66
2	78	77	76
5	72	72	74
7	68	74	73

10

The results in Table 4 show that the activity did not decrease at any of the temperatures of storage between the 1 and 7 day incubation period.

Claims

1. A microorganism which is *Rhodococcus rhodochrous* strain NCIMB 41164 or a mutant thereof.
- 5 2. A method of culturing the microorganism *Rhodococcus rhodochrous* strain NCIMB 41164 or mutant thereof in a culture medium that contains urea or a derivative of urea.
3. A method according to claim 2 in which urea or urea derivative is  
10 introduced into the culture medium at least six hours after the start of growth of the microorganism.
4. A method according to claim 2 or claim 3 in which the culture medium contains less than 0.2 g/l urea or the urea derivative for at least the first 6 hours  
15 of culturing the microorganism and thereafter urea or the urea derivative is added to the culture medium.
5. A method according to any of claims 2 to 4 in which the culture medium contains less than 0.2 g/l urea or the urea derivative for at least the first 12  
20 hours of culturing the microorganism and thereafter urea or the urea derivative is added to the culture medium.
6. A method according to any of claims 2 to 5 in which urea or the urea derivative is added to the culture medium within 48 hours of culturing.
- 25 7. A nitrile hydratase obtainable from a microorganism which is *Rhodococcus rhodochrous* strain NCIMB 41164 or a mutant thereof.
8. A process of preparing an amide from the corresponding nitrile wherein  
30 the nitrile is subjected to a hydration reaction in an aqueous medium in the presence of a biocatalyst selected from the group consisting of a microorganism

which is a *Rhodococcus rhodochrous* strain NCIMB 41164, a mutant thereof and a nitrile hydratase obtainable from *Rhodococcus rhodochrous* strain NCIMB 41164 or a mutant thereof.

- 5 9. A process according to claim 8 in which the amide is (meth)acrylamide.
10. A process according to claim 9 in which the biocatalyst is introduced into an aqueous medium and (meth)acrylonitrile is fed into the aqueous medium such that the concentration of (meth)acrylonitrile in the aqueous medium is  
10 maintained at up to 6% by weight.
11. A process according to claim 10 in which the reaction continues until the concentration of acrylamide is between 30 and 55% by weight.
- 15 12. A process according to any of claims 8 to 11 in which the biocatalyst is recycled and reused.
13. A method of improving the biocatalytic activity of a microorganism, in which the microorganism is cultured in a culture medium that comprises urea or  
20 a derivative of urea ,  
wherein urea or the derivative of urea is introduced into the culture medium at least 6 hours after the start of growth of the microorganism.
14. A method according to claim 13 in which the culture medium contains less  
25 than 0.2 g/l urea or the derivative of urea for at least the first 6 hours of culturing the microorganism and thereafter urea or the derivative of urea is added to the culture medium.
15. A method according to claim 13 or claim 14 in which the culture medium  
30 contains less than 0.2 g/l urea or the derivative of urea for at least the first 12

hours of culturing the microorganism and thereafter urea or the derivative of urea is added to the culture medium.

16. A method according to any of claims 13 to 15 in which urea or the urea  
5 derivative is added to the culture medium within 48 hours of culturing.

17. A method according to any of claims 13 to 16 in which the microorganism is capable of producing a nitrile hydratase.

10 18. A method according to any of claims 13 to 17 in which the microorganism is of the Rhodococcus genus, preferably Rhodococcus rhodochrous species.

19. A process of preparing an amide from the corresponding nitrile wherein the nitrile is subjected to a hydration reaction in an aqueous medium in the  
15 presence of a biocatalyst selected from the group consisting of a microorganism which is capable of producing a nitrile hydratase, wherein the microorganism has been cultured by the method according to any of claims 13 to 18.

20 20. A process according to claim 19 in which the amide is (meth) acrylamide.

21. A process according to claim 19 in which the biocatalyst is introduced into an aqueous medium and (meth)acrylonitrile is fed into the aqueous medium such that the concentration of (meth)acrylonitrile in the aqueous medium is  
25 maintained at up to 6% by weight.

22. A process according to claim 21 in which the reaction continues until the concentration of acrylamide is between 30 and 55% by weight.

30 23. A process according to any of claims 19 to 22 in which the biocatalyst is recycled and reused.

24. An aqueous composition comprising a biocatalyst that is or is obtainable from the microorganism *Rhodococcus rhodochrous* strain NCIMB 41164 or a mutant thereof and wherein the biocatalyst is in the form of a non-actively growing free cell microorganism.

25. A method of storing the biocatalyst that is or is obtainable from the microorganism *Rhodococcus rhodochrous* strain NCIMB 41164 or a mutant thereof in the form of a non-actively growing free cell microorganism, in an aqueous storage medium.

26. A method according to claim 25 in which the biocatalyst is stored at a temperature above its freezing point, preferably above 0°C and more preferably between 4 and 30°C.

27. A method according to claim 25 or claim 26 in which the biocatalyst is stored for a period of at least two days, preferably between 3 and 28 days and more preferably between 5 and 14 days.

28. A composition obtainable by the method according to any of claims 25 to 27.

29. A nitrile hydratase obtainable from the composition according to claim 24 or obtainable by the method according to any of claims 25 to 27.


30. A method of producing an amide by contacting the corresponding nitrile with a nitrile hydratase, wherein the nitrile hydratase is obtainable from a composition according to claim 24 or obtainable by a method according to claims 25 to 27.

31. A method according to claim 30 in which the amide is (meth)acrylamide.

AbstractMicroorganism

A microorganism which is Rhodococcus rhodochrous strain NCIMB 41164 or a mutant thereof. A method of culturing the microorganism in a culture medium comprising urea or urea derivative is claimed. A nitrile hydratase obtainable from the microorganism is claimed. Also claimed is a process of preparing an amide from the corresponding nitrile wherein the nitrile is subjected to a hydration reaction in an aqueous medium in the presence of a biocatalyst selected from the group consisting of a microorganism which is a Rhodococcus rhodochrous strain NCIMB 41164, a mutant thereof and a nitrile hydratase obtainable from Rhodococcus rhodochrous strain NCIMB 41164 or a mutant thereof. Also claimed is a method of storing the Rhodococcus rhodochrous NCIMB 41164.

PCT/EP2004/013252



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**